

Search for Radiative Penguin Decays $B^+ \rightarrow \rho^+ \gamma$, $B^0 \rightarrow \rho^0 \gamma$, and $B^0 \rightarrow \omega \gamma$

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ J.-M. Gaillard,¹ A. Hicheur,¹ Y. Karyotakis,¹ J. P. Lees,¹
V. Tisserand,¹ A. Zghiche,¹ A. Palano,² A. Pompili,² J. C. Chen,³ N. D. Qi,³ G. Rong,³ P. Wang,³ Y. S. Zhu,³
G. Eigen,⁴ I. Ofte,⁴ B. Stugu,⁴ G. S. Abrams,⁵ A. W. Borgland,⁵ A. B. Breon,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵
R. N. Cahn,⁵ E. Charles,⁵ C. T. Day,⁵ M. S. Gill,⁵ A. V. Gritsan,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ R. W. Kadel,⁵
J. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Kukartsev,⁵ G. Lynch,⁵ L. M. Mir,⁵ P. J. Oddone,⁵
T. J. Orimoto,⁵ M. Pripstein,⁵ N. A. Roe,⁵ M. T. Ronan,⁵ V. G. Shelkov,⁵ W. A. Wenzel,⁵ M. Barrett,⁶
K. E. Ford,⁶ T. J. Harrison,⁶ A. J. Hart,⁶ C. M. Hawkes,⁶ S. E. Morgan,⁶ A. T. Watson,⁶ M. Fritsch,⁷ K. Goetzen,⁷
T. Held,⁷ H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ M. Steinke,⁷ J. T. Boyd,⁸ N. Chevalier,⁸ W. N. Cottingham,⁸
M. P. Kelly,⁸ T. E. Latham,⁸ F. F. Wilson,⁸ T. Cuhadar-Donszelmann,⁹ C. Hearty,⁹ N. S. Knecht,⁹ T. S. Mattison,⁹
J. A. McKenna,⁹ D. Thiessen,⁹ A. Khan,¹⁰ P. Kyberd,¹⁰ L. Teodorescu,¹⁰ A. E. Blinov,¹¹ V. E. Blinov,¹¹
V. P. Druzhinin,¹¹ V. B. Golubev,¹¹ V. N. Ivanchenko,¹¹ E. A. Kravchenko,¹¹ A. P. Onuchin,¹¹ S. I. Serebnyakov,¹¹
Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ A. N. Yushkov,¹¹ D. Best,¹² M. Bruinsma,¹² M. Chao,¹² I. Eschrich,¹²
D. Kirkby,¹² A. J. Lankford,¹² M. Mandelkern,¹² R. K. Mommsen,¹² W. Roethel,¹² D. P. Stoker,¹² C. Buchanan,¹³
B. L. Hartfiel,¹³ S. D. Foulkes,¹⁴ J. W. Gary,¹⁴ B. C. Shen,¹⁴ K. Wang,¹⁴ D. del Re,¹⁵ H. K. Hadavand,¹⁵
E. J. Hill,¹⁵ D. B. MacFarlane,¹⁵ H. P. Paar,¹⁵ Sh. Rahatlou,¹⁵ V. Sharma,¹⁵ J. W. Berryhill,¹⁶ C. Campagnari,¹⁶
B. Dahmes,¹⁶ O. Long,¹⁶ A. Lu,¹⁶ M. A. Mazur,¹⁶ J. D. Richman,¹⁶ W. Verkerke,¹⁶ T. W. Beck,¹⁷ A. M. Eisner,¹⁷
C. A. Heusch,¹⁷ J. Kroseberg,¹⁷ W. S. Lockman,¹⁷ G. Nesom,¹⁷ T. Schalk,¹⁷ B. A. Schumm,¹⁷ A. Seiden,¹⁷
P. Spradlin,¹⁷ D. C. Williams,¹⁷ M. G. Wilson,¹⁷ J. Albert,¹⁸ E. Chen,¹⁸ G. P. Dubois-Felsmann,¹⁸
A. Dvoretzskii,¹⁸ D. G. Hitlin,¹⁸ I. Narsky,¹⁸ T. Piatenko,¹⁸ F. C. Porter,¹⁸ A. Ryd,¹⁸ A. Samuel,¹⁸ S. Yang,¹⁸
S. Jayatilake,¹⁹ G. Mancinelli,¹⁹ B. T. Meadows,¹⁹ M. D. Sokoloff,¹⁹ T. Abe,²⁰ F. Blanc,²⁰ P. Bloom,²⁰
S. Chen,²⁰ W. T. Ford,²⁰ U. Nauenberg,²⁰ A. Olivas,²⁰ P. Rankin,²⁰ J. G. Smith,²⁰ J. Zhang,²⁰ L. Zhang,²⁰
A. Chen,²¹ J. L. Harton,²¹ A. Soffer,²¹ W. H. Toki,²¹ R. J. Wilson,²¹ Q. L. Zeng,²¹ D. Altenburg,²² T. Brandt,²²
J. Brose,²² M. Dickopp,²² E. Feltresi,²² A. Hauke,²² H. M. Lacker,²² R. Müller-Pfefferkorn,²² R. Nogowski,²²
S. Otto,²² A. Petzold,²² J. Schubert,²² K. R. Schubert,²² R. Schwierz,²² B. Spaan,²² J. E. Sundermann,²²
D. Bernard,²³ G. R. Bonneaud,²³ F. Brochard,²³ P. Grenier,²³ S. Schrenk,²³ Ch. Thiebaux,²³ G. Vasileiadis,²³
M. Verderi,²³ D. J. Bard,²⁴ P. J. Clark,²⁴ D. Lavin,²⁴ F. Muheim,²⁴ S. Playfer,²⁴ Y. Xie,²⁴ M. Andreotti,²⁵
V. Azzolini,²⁵ D. Bettoni,²⁵ C. Bozzi,²⁵ R. Calabrese,²⁵ G. Cibinetto,²⁵ E. Luppi,²⁵ M. Negrini,²⁵ L. Piemontese,²⁵
A. Sarti,²⁵ E. Treadwell,²⁶ F. Anulli,²⁷ R. Baldini-Ferrolì,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ G. Finocchiaro,²⁷
P. Patteri,²⁷ I. M. Peruzzi,²⁷ M. Piccolo,²⁷ A. Zallo,²⁷ A. Buzzo,²⁸ R. Capra,²⁸ R. Contri,²⁸ G. Crosetti,²⁸
M. Lo Vetere,²⁸ M. Macri,²⁸ M. R. Monge,²⁸ S. Passaggio,²⁸ C. Patrignani,²⁸ E. Robutti,²⁸ A. Santroni,²⁸
S. Tosi,²⁸ S. Bailey,²⁹ G. Brandenburg,²⁹ K. S. Chaisanguanthum,²⁹ M. Morii,²⁹ E. Won,²⁹ R. S. Dubitzky,³⁰
U. Langenegger,³⁰ W. Bhimji,³¹ D. A. Bowerman,³¹ P. D. Dauncey,³¹ U. Egede,³¹ J. R. Gaillard,³¹ G. W. Morton,³¹
J. A. Nash,³¹ M. B. Nikolich,³¹ G. P. Taylor,³¹ M. J. Charles,³² G. J. Grenier,³² U. Mallik,³² J. Cochran,³³
H. B. Crawley,³³ J. Lamsa,³³ W. T. Meyer,³³ S. Prell,³³ E. I. Rosenberg,³³ A. E. Rubin,³³ J. Yi,³³ M. Biasini,³⁴
R. Covarelli,³⁴ M. Pippi,³⁴ M. Davier,³⁵ X. Giroux,³⁵ G. Grosdidier,³⁵ A. Höcker,³⁵ S. Laplace,³⁵ F. Le Diberder,³⁵
V. Lepeltier,³⁵ A. M. Lutz,³⁵ T. C. Petersen,³⁵ S. Plaszczynski,³⁵ M. H. Schune,³⁵ L. Tantot,³⁵ G. Wormser,³⁵
C. H. Cheng,³⁶ D. J. Lange,³⁶ M. C. Simani,³⁶ D. M. Wright,³⁶ A. J. Bevan,³⁷ C. A. Chavez,³⁷ J. P. Coleman,³⁷
I. J. Forster,³⁷ J. R. Fry,³⁷ E. Gabathuler,³⁷ R. Gamet,³⁷ D. E. Hutchcroft,³⁷ R. J. Parry,³⁷ D. J. Payne,³⁷
R. J. Sloane,³⁷ C. Touramanis,³⁷ J. J. Back,^{38,*} C. M. Cormack,³⁸ P. F. Harrison,^{38,*} F. Di Lodovico,³⁸
G. B. Mohanty,^{38,*} C. L. Brown,³⁹ G. Cowan,³⁹ R. L. Flack,³⁹ H. U. Flaecher,³⁹ M. G. Green,³⁹ P. S. Jackson,³⁹
T. R. McMahon,³⁹ S. Ricciardi,³⁹ F. Salvatore,³⁹ M. A. Winter,³⁹ D. Brown,⁴⁰ C. L. Davis,⁴⁰ J. Allison,⁴¹
N. R. Barlow,⁴¹ R. J. Barlow,⁴¹ P. A. Hart,⁴¹ M. C. Hodgkinson,⁴¹ G. D. Lafferty,⁴¹ A. J. Lyon,⁴¹ J. C. Williams,⁴¹
A. Farbin,⁴² W. D. Hulsbergen,⁴² A. Jawahery,⁴² D. Kovalskyi,⁴² C. K. Lae,⁴² V. Lillard,⁴² D. A. Roberts,⁴²
G. Blaylock,⁴³ C. Dallapiccola,⁴³ K. T. Flood,⁴³ S. S. Hertzbach,⁴³ R. Kofler,⁴³ V. B. Koptchev,⁴³ T. B. Moore,⁴³
S. Saremi,⁴³ H. Staengle,⁴³ S. Willocq,⁴³ R. Cowan,⁴⁴ G. Sciolla,⁴⁴ S. J. Sekula,⁴⁴ F. Taylor,⁴⁴ R. K. Yamamoto,⁴⁴
D. J. J. Mangeol,⁴⁵ P. M. Patel,⁴⁵ S. H. Robertson,⁴⁵ A. Lazzaro,⁴⁶ V. Lombardo,⁴⁶ F. Palombo,⁴⁶
J. M. Bauer,⁴⁷ L. Cremaldi,⁴⁷ V. Eschenburg,⁴⁷ R. Godang,⁴⁷ R. Kroeger,⁴⁷ J. Reidy,⁴⁷ D. A. Sanders,⁴⁷

D. J. Summers,⁴⁷ H. W. Zhao,⁴⁷ S. Brunet,⁴⁸ D. Côté,⁴⁸ P. Taras,⁴⁸ H. Nicholson,⁴⁹ N. Cavallo,^{50,†} F. Fabozzi,^{50,†} C. Gatto,⁵⁰ L. Lista,⁵⁰ D. Monorchio,⁵⁰ P. Paolucci,⁵⁰ D. Piccolo,⁵⁰ C. Sciacca,⁵⁰ M. Baak,⁵¹ H. Bulten,⁵¹ G. Raven,⁵¹ H. L. Snoek,⁵¹ L. Wilden,⁵¹ C. P. Jessop,⁵² J. M. LoSecco,⁵² T. Allmendinger,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ E. Torrence,⁵⁴ F. Colechia,⁵⁵ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵ M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ G. Tiozzo,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ Ch. de la Vaissière,⁵⁶ L. Del Buono,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Malcles,⁵⁶ J. Ocariz,⁵⁶ M. Pivk,⁵⁶ L. Roos,⁵⁶ S. T'Jampens,⁵⁶ G. Therin,⁵⁶ P. F. Manfredi,⁵⁷ V. Re,⁵⁷ P. K. Behera,⁵⁸ L. Gladney,⁵⁸ Q. H. Guo,⁵⁸ J. Panetta,⁵⁸ C. Angelini,⁵⁹ G. Batignani,⁵⁹ S. Bettarini,⁵⁹ M. Bondioli,⁵⁹ F. Bucci,⁵⁹ G. Calderini,⁵⁹ M. Carpinelli,⁵⁹ F. Forti,⁵⁹ M. A. Giorgi,⁵⁹ A. Lusiani,⁵⁹ G. Marchiori,⁵⁹ F. Martinez-Vidal,^{59,‡} M. Morganti,⁵⁹ N. Neri,⁵⁹ E. Paoloni,⁵⁹ M. Rama,⁵⁹ G. Rizzo,⁵⁹ F. Sandrelli,⁵⁹ J. Walsh,⁵⁹ M. Haire,⁶⁰ D. Judd,⁶⁰ K. Paick,⁶⁰ D. E. Wagoner,⁶⁰ N. Danielson,⁶¹ P. Elmer,⁶¹ Y. P. Lau,⁶¹ C. Lu,⁶¹ V. Miftakov,⁶¹ J. Olsen,⁶¹ A. J. S. Smith,⁶¹ A. V. Telnov,⁶¹ F. Bellini,⁶² G. Cavoto,^{61,62} R. Faccini,⁶² F. Ferrarotto,⁶² F. Ferroni,⁶² M. Gaspero,⁶² L. Li Gioi,⁶² M. A. Mazzoni,⁶² S. Morganti,⁶² M. Pierini,⁶² G. Piredda,⁶² F. Safai Tehrani,⁶² C. Voena,⁶² S. Christ,⁶³ G. Wagner,⁶³ R. Waldi,⁶³ T. Adye,⁶⁴ N. De Groot,⁶⁴ B. Franek,⁶⁴ N. I. Geddes,⁶⁴ G. P. Gopal,⁶⁴ E. O. Olaiya,⁶⁴ R. Aleksan,⁶⁵ S. Emery,⁶⁵ A. Gaidot,⁶⁵ S. F. Ganzhur,⁶⁵ P.-F. Giraud,⁶⁵ G. Hamel de Monchenault,⁶⁵ W. Kozanecki,⁶⁵ M. Legendre,⁶⁵ G. W. London,⁶⁵ B. Mayer,⁶⁵ G. Schott,⁶⁵ G. Vasseur,⁶⁵ Ch. Yèche,⁶⁵ M. Zito,⁶⁵ M. V. Purohit,⁶⁶ A. W. Weidemann,⁶⁶ J. R. Wilson,⁶⁶ F. X. Yumiceva,⁶⁶ D. Aston,⁶⁷ R. Bartoldus,⁶⁷ N. Berger,⁶⁷ A. M. Boyarski,⁶⁷ O. L. Buchmueller,⁶⁷ R. Claus,⁶⁷ M. R. Convery,⁶⁷ M. Cristinziani,⁶⁷ G. De Nardo,⁶⁷ D. Dong,⁶⁷ J. Dorfan,⁶⁷ D. Dujmic,⁶⁷ W. Dunwoodie,⁶⁷ E. E. Elsen,⁶⁷ S. Fan,⁶⁷ R. C. Field,⁶⁷ T. Glanzman,⁶⁷ S. J. Gowdy,⁶⁷ T. Hadig,⁶⁷ V. Halyo,⁶⁷ C. Hast,⁶⁷ T. Hryn'ova,⁶⁷ W. R. Innes,⁶⁷ M. H. Kelsey,⁶⁷ P. Kim,⁶⁷ M. L. Kocian,⁶⁷ D. W. G. S. Leith,⁶⁷ J. Libby,⁶⁷ S. Luitz,⁶⁷ V. Luth,⁶⁷ H. L. Lynch,⁶⁷ H. Marsiske,⁶⁷ R. Messner,⁶⁷ D. R. Muller,⁶⁷ C. P. O'Grady,⁶⁷ V. E. Ozcan,⁶⁷ A. Perazzo,⁶⁷ M. Perl,⁶⁷ S. Petrak,⁶⁷ B. N. Ratcliff,⁶⁷ A. Roodman,⁶⁷ A. A. Salnikov,⁶⁷ R. H. Schindler,⁶⁷ J. Schwiening,⁶⁷ G. Simi,⁶⁷ A. Snyder,⁶⁷ A. Soha,⁶⁷ J. Stelzer,⁶⁷ D. Su,⁶⁷ M. K. Sullivan,⁶⁷ J. Va'vra,⁶⁷ S. R. Wagner,⁶⁷ M. Weaver,⁶⁷ A. J. R. Weinstein,⁶⁷ W. J. Wisniewski,⁶⁷ M. Wittgen,⁶⁷ D. H. Wright,⁶⁷ A. K. Yarritu,⁶⁷ C. C. Young,⁶⁷ P. R. Burchat,⁶⁸ A. J. Edwards,⁶⁸ T. I. Meyer,⁶⁸ B. A. Petersen,⁶⁸ C. Roat,⁶⁸ S. Ahmed,⁶⁹ M. S. Alam,⁶⁹ J. A. Ernst,⁶⁹ M. A. Saeed,⁶⁹ M. Saleem,⁶⁹ F. R. Wappler,⁶⁹ W. Bugg,⁷⁰ M. Krishnamurthy,⁷⁰ S. M. Spanier,⁷⁰ R. Eckmann,⁷¹ H. Kim,⁷¹ J. L. Ritchie,⁷¹ A. Satpathy,⁷¹ R. F. Schwitters,⁷¹ J. M. Izen,⁷² I. Kitayama,⁷² X. C. Lou,⁷² S. Ye,⁷² F. Bianchi,⁷³ M. Bona,⁷³ F. Gallo,⁷³ D. Gamba,⁷³ L. Bosisio,⁷⁴ C. Cartaro,⁷⁴ F. Cossutti,⁷⁴ G. Della Ricca,⁷⁴ S. Dittongo,⁷⁴ S. Grancagnolo,⁷⁴ L. Lanceri,⁷⁴ P. Poropat,^{74,§} L. Vitale,⁷⁴ G. Vuagnin,⁷⁴ R. S. Panvini,⁷⁵ Sw. Banerjee,⁷⁶ C. M. Brown,⁷⁶ D. Fortin,⁷⁶ P. D. Jackson,⁷⁶ R. Kowalewski,⁷⁶ J. M. Roney,⁷⁶ R. J. Sobie,⁷⁶ H. R. Band,⁷⁷ B. Cheng,⁷⁷ S. Dasu,⁷⁷ M. Datta,⁷⁷ A. M. Eichenbaum,⁷⁷ M. Graham,⁷⁷ J. J. Hollar,⁷⁷ J. R. Johnson,⁷⁷ P. E. Kutter,⁷⁷ H. Li,⁷⁷ R. Liu,⁷⁷ A. Mihalyi,⁷⁷ A. K. Mohapatra,⁷⁷ Y. Pan,⁷⁷ R. Prepost,⁷⁷ P. Tan,⁷⁷ J. H. von Wimmersperg-Toeller,⁷⁷ J. Wu,⁷⁷ S. L. Wu,⁷⁷ Z. Yu,⁷⁷ M. G. Greene,⁷⁸ and H. Neal⁷⁸

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

³Institute of High Energy Physics, Beijing 100039, China

⁴University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, CA 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, BC, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, CA 92697, USA

¹³University of California at Los Angeles, Los Angeles, CA 90024, USA

¹⁴University of California at Riverside, Riverside, CA 92521, USA

¹⁵University of California at San Diego, La Jolla, CA 92093, USA

¹⁶University of California at Santa Barbara, Santa Barbara, CA 93106, USA

¹⁷University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, CA 95064, USA

¹⁸California Institute of Technology, Pasadena, CA 91125, USA

- ¹⁹University of Cincinnati, Cincinnati, OH 45221, USA
²⁰University of Colorado, Boulder, CO 80309, USA
²¹Colorado State University, Fort Collins, CO 80523, USA
²²Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany
²³Ecole Polytechnique, LLR, F-91128 Palaiseau, France
²⁴University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
²⁵Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
²⁶Florida A&M University, Tallahassee, FL 32307, USA
²⁷Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
²⁸Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
²⁹Harvard University, Cambridge, MA 02138, USA
³⁰Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
³¹Imperial College London, London, SW7 2AZ, United Kingdom
³²University of Iowa, Iowa City, IA 52242, USA
³³Iowa State University, Ames, IA 50011-3160, USA
³⁴Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
³⁵Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
³⁶Lawrence Livermore National Laboratory, Livermore, CA 94550, USA
³⁷University of Liverpool, Liverpool L69 7ZE, United Kingdom
³⁸Queen Mary, University of London, E1 4NS, United Kingdom
³⁹University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
⁴⁰University of Louisville, Louisville, KY 40292, USA
⁴¹University of Manchester, Manchester M13 9PL, United Kingdom
⁴²University of Maryland, College Park, MD 20742, USA
⁴³University of Massachusetts, Amherst, MA 01003, USA
⁴⁴Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, MA 02139, USA
⁴⁵McGill University, Montréal, QC, Canada H3A 2T8
⁴⁶Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
⁴⁷University of Mississippi, University, MS 38677, USA
⁴⁸Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, QC, Canada H3C 3J7
⁴⁹Mount Holyoke College, South Hadley, MA 01075, USA
⁵⁰Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
⁵¹NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
⁵²University of Notre Dame, Notre Dame, IN 46556, USA
⁵³Ohio State University, Columbus, OH 43210, USA
⁵⁴University of Oregon, Eugene, OR 97403, USA
⁵⁵Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
⁵⁶Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France
⁵⁷Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy
⁵⁸University of Pennsylvania, Philadelphia, PA 19104, USA
⁵⁹Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
⁶⁰Prairie View A&M University, Prairie View, TX 77446, USA
⁶¹Princeton University, Princeton, NJ 08544, USA
⁶²Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
⁶³Universität Rostock, D-18051 Rostock, Germany
⁶⁴Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
⁶⁵DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
⁶⁶University of South Carolina, Columbia, SC 29208, USA
⁶⁷Stanford Linear Accelerator Center, Stanford, CA 94309, USA
⁶⁸Stanford University, Stanford, CA 94305-4060, USA
⁶⁹State University of New York, Albany, NY 12222, USA
⁷⁰University of Tennessee, Knoxville, TN 37996, USA
⁷¹University of Texas at Austin, Austin, TX 78712, USA
⁷²University of Texas at Dallas, Richardson, TX 75083, USA
⁷³Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
⁷⁴Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
⁷⁵Vanderbilt University, Nashville, TN 37235, USA
⁷⁶University of Victoria, Victoria, BC, Canada V8W 3P6
⁷⁷University of Wisconsin, Madison, WI 53706, USA
⁷⁸Yale University, New Haven, CT 06511, USA

(Dated: February 7, 2008)

A search for the decays $B \rightarrow \rho(770)\gamma$ and $B^0 \rightarrow \omega(782)\gamma$ is performed on a sample of 211 million $\Upsilon(4S) \rightarrow B\bar{B}$ events collected by the BABAR detector at the PEP-II asymmetric-energy e^+e^- storage

ring. No evidence for the decays is seen. We set the following limits on the individual branching fractions $\mathcal{B}(B^+ \rightarrow \rho^+\gamma) < 1.8 \times 10^{-6}$, $\mathcal{B}(B^0 \rightarrow \rho^0\gamma) < 0.4 \times 10^{-6}$, and $\mathcal{B}(B^0 \rightarrow \omega\gamma) < 1.0 \times 10^{-6}$ at the 90% confidence level (C.L.). We use the quark model to limit the combined branching fraction $\overline{\mathcal{B}}[B \rightarrow (\rho/\omega)\gamma] < 1.2 \times 10^{-6}$, from which we determine a constraint on the ratio of CKM matrix elements $|V_{td}|/|V_{ts}|$.

PACS numbers: 12.15.Hh, 13.25.Hw

Within the Standard Model (SM), the decays $B \rightarrow \rho\gamma$ and $B^0 \rightarrow \omega\gamma$ proceed primarily through a $b \rightarrow d\gamma$ electromagnetic penguin process that contains a top quark within the loop [1]. The rates for $B^+ \rightarrow \rho^+\gamma$, $B^0 \rightarrow \rho^0\gamma$, and $B^0 \rightarrow \omega\gamma$ [2] are related by the spectator-quark model, and we define the average branching fraction [3], $\overline{\mathcal{B}}[B \rightarrow (\rho/\omega)\gamma] = \frac{1}{2} \left\{ \mathcal{B}(B^+ \rightarrow \rho^+\gamma) + \frac{\tau_{B^+}}{\tau_{B^0}} [\mathcal{B}(B^0 \rightarrow \rho^0\gamma) + \mathcal{B}(B^0 \rightarrow \omega\gamma)] \right\}$, where $\frac{\tau_{B^+}}{\tau_{B^0}}$ is the ratio of B -meson lifetimes. Recent calculations of $\overline{\mathcal{B}}[B \rightarrow (\rho/\omega)\gamma]$ in the SM indicate a range of $(0.9 - 1.8) \times 10^{-6}$ [3, 4]. There may also be contributions resulting from physics beyond the SM [5]. The ratio between the branching fractions for $B \rightarrow (\rho/\omega)\gamma$ and $B \rightarrow K^*\gamma$ is related in the SM to the ratio of Cabibbo-Kobayashi-Maskawa (CKM) matrix elements $|V_{td}|/|V_{ts}|$ [3, 6]. Previous searches by BABAR [7] and CLEO [8] have found no evidence for $B \rightarrow (\rho/\omega)\gamma$ decays.

We search for $B \rightarrow \rho\gamma$ and $B^0 \rightarrow \omega\gamma$ decays in a data sample containing 211 ± 2 million $\Upsilon(4S) \rightarrow B\overline{B}$ decays, collected by the BABAR detector [9] at the PEP-II asymmetric-energy e^+e^- storage ring. The data correspond to an integrated luminosity of 191 fb^{-1} .

The decay $B \rightarrow \rho\gamma$ is reconstructed with $\rho^0 \rightarrow \pi^+\pi^-$ and $\rho^+ \rightarrow \pi^+\pi^0$, while $B^0 \rightarrow \omega\gamma$ is reconstructed with $\omega \rightarrow \pi^+\pi^-\pi^0$. Background comes primarily from $e^+e^- \rightarrow q\overline{q}$ continuum events, where $q = u, d, s, c$, in which a high-energy photon is produced through $\pi^0/\eta \rightarrow \gamma\gamma$ decays or via initial-state radiation (ISR). There are also significant $B\overline{B}$ backgrounds: $B \rightarrow K^*\gamma$, $K^* \rightarrow K\pi$, where a K^\pm is misidentified as a π^\pm ; $B \rightarrow (\rho/\omega)\pi^0$ and $B \rightarrow (\rho/\omega)\eta$, where a high-energy photon comes from the π^0 or η decay; and combinatorial background, mostly from high multiplicity $b \rightarrow s\gamma$ decays.

We select π^\pm candidates from tracks with a momentum transverse to the beam direction greater than 100 MeV/c. The π^\pm selection algorithm combines measurements of energy loss in the tracking system with any associated Cherenkov photons measured by the ring imaging Cherenkov detector. The algorithm is optimized to reduce backgrounds from K^\pm produced in $b \rightarrow s\gamma$ processes [7].

Neutral pion candidates are identified as pairs of neutral energy-deposits reconstructed in the CsI crystal calorimeter, each with an energy greater than 50 MeV in the laboratory frame. For $B^0 \rightarrow \omega\gamma$ ($B^+ \rightarrow \rho^+\gamma$) decays, the invariant mass of the pair is required to satisfy

$110 < m_{\gamma\gamma} < 150 \text{ MeV}/c^2$ ($117 < m_{\gamma\gamma} < 145 \text{ MeV}/c^2$). To reduce combinatorial background, we require the cosine of the opening angle between the daughter photons in the laboratory frame be greater than 0.6; this selection retains 98% of π^0 from signal decays.

A ρ^0 candidate is reconstructed by selecting two identified pions that have opposite charge and a common vertex. We obtain ρ^+ candidates by pairing π^0 candidates with an identified π^+ . The ω candidates are reconstructed by combining a π^0 candidate with pairs of oppositely charged pion candidates that originate from a common vertex; the charged pion pair must be consistent with originating from the interaction region to suppress K_s^0 decays. We select ρ (ω) candidates with an invariant mass satisfying $630 < m_{\pi\pi} < 940 \text{ MeV}/c^2$ ($764 < m_{\pi^+\pi^-\pi^0} < 795 \text{ MeV}/c^2$).

The high-energy photon from the signal B decay is identified as a neutral energy-deposit in the calorimeter. We require that the deposit meets a number of criteria designed to eliminate background from charged particles and hadronic showers [10]. We veto photons from $\pi^0(\eta)$ decay by requiring that the invariant mass of the candidate combined with any other photon of laboratory energy greater than 30 (250) MeV not to be within the range 105 to 155 MeV/ c^2 (500 to 590 MeV/ c^2).

The photon and ρ/ω candidates are combined to form the B -meson candidates. We define $\Delta E^* \equiv E_B^* - E_{\text{beam}}^*$, where E_B^* is the center-of-mass (CM) energy of the B -meson candidate and E_{beam}^* is the CM beam energy. The ΔE^* distribution of Monte Carlo (MC) simulated signal events is centered at zero, with a resolution of about 50 MeV. We also define the beam-energy-substituted mass $m_{\text{ES}} \equiv \sqrt{E_{\text{beam}}^{*2} - \mathbf{p}_B^{*2}}$, where \mathbf{p}_B^* is the CM momentum of the B candidate. Signal MC events peak in m_{ES} at the mass of the B -meson, m_B , with a resolution of 3 MeV/ c^2 . The distribution of continuum and combinatorial $B\overline{B}$ background peaks in neither m_{ES} nor ΔE^* ; the background distributions of $B \rightarrow K^*\gamma$, $B \rightarrow (\rho/\omega)\pi^0$ and $B \rightarrow (\rho/\omega)\eta$ peak at m_B in m_{ES} and between -190 MeV and -60 MeV in ΔE^* . We consider candidates in the ranges $-0.3 < \Delta E^* < 0.3 \text{ GeV}$ and $5.20 < m_{\text{ES}} < 5.29 \text{ GeV}/c^2$ to incorporate sidebands that allow the combinatorial background yields to be extracted from a fit to the data.

Several variables that distinguish between signal and continuum events are combined in a neural network [11]. The input variables depend mainly on the rest of the event (ROE), defined to be all charged tracks and neu-

tral energy deposits in the calorimeter not used to reconstruct the B candidate. To reject ISR events, we compute the ratio of second-to-zeroth order Fox-Wolfram moments [12] for the ROE and the ρ/ω candidate, in the frame recoiling against the photon momentum. To discriminate between the jet-like continuum background and the more spherically-symmetric signal events, we compute the angle between the photon and the thrust axis of the ROE in the CM frame and the moments $L_i \equiv \sum_j p_j^* \cdot |\cos \theta_j^*|^i / \sum_j p_j^*$, where p_j^* and θ_j^* are the momentum and angle with respect to an axis, respectively, for each particle j in the ROE. We use L_1 , L_2 , and L_3 with respect to the thrust axis of the ROE, as well as L_1 with respect to the photon direction. Differences in lepton and kaon production between background and B decays are exploited by including *BABAR* flavor tagging variables [13] as well as the maximum CM momentum and number of K^\pm and K_s^0 in the ROE. For the $B^0 \rightarrow (\rho^0/\omega)\gamma$ modes, we also use the separation along the beam axis of the B -meson candidate and ROE vertices; to remove poorly reconstructed events we require the separation be less than 4 mm. A separate neural network is trained for each mode. We make a loose selection on the output of the neural network, \mathcal{N} , that retains around 80% of the signal events.

To suppress background, we combine a number of signal-decay variables in a Fisher discriminant [14], \mathcal{F} , separately for each mode. We calculate the B -meson production angle θ_B^* , the ρ/ω helicity angle θ_H , which is defined with respect to the normal of the decay plane for ω candidates, and the ω Dalitz angle θ_D [7]. To reject $B \rightarrow \rho(\pi^0/\eta)$ and $B \rightarrow \omega(\pi^0/\eta)$ events in the $B^+ \rightarrow \rho^+\gamma$ and $B^0 \rightarrow \omega\gamma$ ($B^0 \rightarrow \rho^0\gamma$) selection, we require $|\cos \theta_H| < 0.70$ (0.75).

After applying the \mathcal{N} and $|\cos \theta_H|$ criteria the expected average candidate multiplicity in signal events is 1.15, 1.03 and 1.14 for $B^+ \rightarrow \rho^+\gamma$, $B^0 \rightarrow \rho^0\gamma$ and $B^0 \rightarrow \omega\gamma$, respectively; in events with multiple candidates the one with the smallest value of $|\Delta E^*|$ is retained.

The signal yield is determined from an extended maximum likelihood fit to the selected data. We fit the four-dimensional distribution of m_{ES} , ΔE^* , \mathcal{F} and \mathcal{N} . For the $B \rightarrow \rho\gamma$ fits, five event hypotheses are considered: signal, continuum background, combinatorial B -background, peaking $B \rightarrow \rho(\pi^0/\eta)$ background and peaking $B \rightarrow K^*\gamma$ background. For the $B^0 \rightarrow \omega\gamma$ fit we consider only signal, continuum background, and peaking $B \rightarrow \omega(\pi^0/\eta)$ background. The correlations among the observables are small; therefore, we assume that the probability density function (PDF) $\mathcal{P}(\vec{x}_j; \vec{\alpha}_i)$ for each hypothesis is the product of individual PDFs for the variables $\vec{x}_j = \{m_{\text{ES}}, \Delta E^*, \mathcal{F}, \mathcal{N}\}$ given the set of parameters $\vec{\alpha}_i$. The likelihood function is a product over all N_k can-

didate events of the sum of the PDFs,

$$\mathcal{L}_k = \exp \left(- \sum_{i=1}^{N_{\text{hyp}}} n_i \right) \cdot \left[\prod_{j=1}^{N_k} \left(\sum_{i=1}^{N_{\text{hyp}}} n_i \mathcal{P}_i(\vec{x}_j; \vec{\alpha}_i) \right) \right],$$

where n_i is the yield of each hypothesis, k is the $B \rightarrow (\rho/\omega)\gamma$ mode, and $N_{\text{hyp}} = 5(3)$ for $B \rightarrow \rho\gamma$ ($B \rightarrow \omega\gamma$).

The m_{ES} and ΔE^* PDFs are parameterized by a Crystal Ball function [15] for both the signal and peaking background. The parameterization is determined from signal MC samples, except the mean of the ΔE^* distribution, which is offset by the observed difference between data and MC samples of $B \rightarrow K^*\gamma$ decays. The continuum background m_{ES} and ΔE^* distributions are parameterized by an ARGUS threshold function [16] and a second-order polynomial, respectively. The combinatorial B background is described by a smoothed distribution [17] determined from MC events in both m_{ES} and ΔE^* . The distribution of \mathcal{N} for signal and $B\bar{B}$ background is parameterized by a Crystal Ball function. The \mathcal{N} distribution for continuum is determined from sideband data, and a histogram is used as the PDF. The distribution of \mathcal{F} is parameterized by smoothed histograms of sideband data for the continuum background and MC events for all other hypotheses.

The fit to the data determines the shape parameters of the continuum background m_{ES} and ΔE^* PDFs, as well as the signal, continuum background and combinatorial $B\bar{B}$ background yields. All other parameters are fixed, including the peaking $B\bar{B}$ background yields. A combined fit is also performed relating the modes using the definition of $\bar{B}[B \rightarrow (\rho/\omega)\gamma]$ to determine an effective yield (n_{eff}) assuming $n(B^+ \rightarrow \rho^+\gamma) = n_{\text{eff}} \cdot \epsilon(B^+ \rightarrow \rho^+\gamma)$ and $n(B^0 \rightarrow (\rho^0/\omega)\gamma) = \frac{1}{2} \frac{\tau_{B^0}}{\tau_{B^+}} n_{\text{eff}} \cdot \epsilon(B^0 \rightarrow (\rho^0/\omega)\gamma)$ where n and ϵ are the yields and reconstruction efficiencies of each mode; the efficiencies include the daughter branching fractions. We take $\frac{\tau_{B^0}}{\tau_{B^+}} = 1.086 \pm 0.017$ [18]. Fig. 1 shows the projections of the combined fit results compared to the data. The results for the individual mode signal yields and n_{eff} are given in Table I. The significance is computed as $\sqrt{2\Delta \log \mathcal{L}}$ where $\Delta \log \mathcal{L}$ is the log likelihood difference between the best fit and the null-signal hypothesis. No significant signal is observed.

The most important systematic uncertainties are associated with the modeling of $B\bar{B}$ backgrounds, the fixed parameters of the PDFs used in the fit, and the signal reconstruction efficiency. The first two contribute to the uncertainties on the signal yields.

The uncertainty on the peaking $B \rightarrow K^*\gamma$ background is dominated by the K^\pm misidentification rate; the rate is corrected by the difference in K^\pm misidentification between data and MC samples of D^* decays, with the whole correction taken as the uncertainty. For the $B^+ \rightarrow \rho^+(\pi^0/\eta)$, $B^0 \rightarrow \rho^0\eta$ and $B^0 \rightarrow \omega(\pi^0/\eta)$ peaking background decays, we vary the branching fractions by either one standard deviation from the measured

TABLE I: The signal yield (n_{sig}), continuum background yield (n_{cont}), peaking background (n_{peak}), significance in standard deviations σ , efficiency (ϵ), and branching fraction (\mathcal{B}) central value and upper limit at the 90% C.L for each mode. The results of the combined fit are shown in the bottom row where n_{sig} is equal to n_{eff} , which is described in the text. When two errors are quoted, the first is statistical and the second is systematic.

Mode	n_{sig}	n_{cont}	n_{peak}	Significance (σ)	$\epsilon(\%)$	$\mathcal{B}(10^{-6})$	$\mathcal{B}(10^{-6})$ 90% C.L.
$B^+ \rightarrow \rho^+ \gamma$	26^{+15+2}_{-14-2}	6850 ± 90	18 ± 4	1.9	13.2 ± 1.4	$0.9^{+0.6}_{-0.5} \pm 0.1$	< 1.8
$B^0 \rightarrow \rho^0 \gamma$	$0.3^{+7.2+1.7}_{-5.4-1.6}$	4269 ± 73	18 ± 7	0.0	15.8 ± 1.9	$0.0 \pm 0.2 \pm 0.1$	< 0.4
$B^0 \rightarrow \omega \gamma$	$8.3^{+5.7+1.3}_{-4.5-1.9}$	1378 ± 37	$2.6^{+0.8}_{-1.2}$	1.5	8.6 ± 0.9	$0.5 \pm 0.3 \pm 0.1$	< 1.0
Combined	$269^{+126+40}_{-120-45}$	—	—	2.1	—	$0.6 \pm 0.3 \pm 0.1$	< 1.2




FIG. 1: Projections of the combined fit to $B \rightarrow \rho \gamma$ and $B^0 \rightarrow \omega \gamma$ in the four discriminating variables: (a) m_{ES} , (b) ΔE^* , (c) \mathcal{N} , and (d) \mathcal{F} . The points are data, the solid line is the total PDF and the dashed line is the background only PDF. The selections applied, unless the variable is projected, are: $5.272 < m_{\text{ES}} < 5.286 \text{ GeV}/c^2$, $-0.10 < \Delta E^* < 0.05 \text{ GeV}$ and $\mathcal{N} > 0.9$; the selection efficiencies for signal events are 45%, 57%, 70%, and 44% for the m_{ES} , ΔE^* , \mathcal{N} and \mathcal{F} projections, respectively.

values or between zero and the measured upper limit if the decay has not been observed [19, 20]; the value of the $B^0 \rightarrow \rho^0 \pi^0$ branching fraction is varied between zero and 5.1×10^{-6} [20, 21]. The uncertainty on the peaking background of each mode is shown in Table I. We find the bias from neglecting the $B \rightarrow K^* \gamma$ background and combinatorial $B\bar{B}$ background in the fit to $B^0 \rightarrow \omega \gamma$ candidates is $1.1^{+1.9}_{-1.1}$ events; the corrected yield is given in Table I. To estimate the uncertainty related to the extraction of the signal m_{ES} and ΔE^* PDFs from MC distributions, we vary the parameters within their errors. The variation in the fitted signal yield is taken as a systematic uncertainty. The uncertainty related to the statistics of the histogram PDF that describes the continuum \mathcal{N} distribution is evaluated by varying the binning and by using a

fifth-order polynomial as an alternative PDF. Several different control samples of data and MC events were used to determine alternative PDFs for the different hypotheses; none of these resulted in a significant change to the fitted signal yield.

The signal efficiency systematic error contains uncertainties from tracking, particle identification, photon/ π^0 reconstruction, photon selection and the neural network selection that are determined as in Ref. [22]. We determine the effect of correlations among the fit variables by using an ensemble of MC experiments of parameterized continuum background simulations embedded in samples of fully simulated signal and $B\bar{B}$ background events. No bias is observed within the statistical error on the mean yields from this ensemble, which is taken as a multiplicative systematic uncertainty. The total multiplicative systematic error values are 11%, 13% and 10% for $B^+ \rightarrow \rho^+ \gamma$, $B^0 \rightarrow \rho^0 \gamma$ and $B^0 \rightarrow \omega \gamma$, respectively. The corrected signal efficiencies and their uncertainties are shown in Table I.

In calculating branching fractions, we assume $\mathcal{B}(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = \mathcal{B}(\Upsilon(4S) \rightarrow B^+ B^-) = 0.5$. The 90% C.L. is taken as the largest value of the efficiency-corrected signal yield at which $2\Delta \log \mathcal{L} = 1.28^2$. We include systematic uncertainties by increasing the efficiency corrected signal yield by 1.28 times its systematic uncertainty. Table I shows the resulting upper limits on the branching fractions.

Using the measured value of $\mathcal{B}(B \rightarrow K^* \gamma)$ [22], we calculate a limit of $\bar{\mathcal{B}}[B \rightarrow (\rho/\omega) \gamma] / \mathcal{B}(B \rightarrow K^* \gamma) < 0.029$ at 90% C.L. This limit is used to constrain the ratio of CKM elements $|V_{td}/V_{ts}|$ by means of the equation [3, 6]:

$$\frac{\bar{\mathcal{B}}[B \rightarrow (\rho/\omega) \gamma]}{\mathcal{B}(B \rightarrow K^* \gamma)} = \left| \frac{V_{td}}{V_{ts}} \right|^2 \left(\frac{1 - m_\rho^2/M_B^2}{1 - m_{K^*}^2/M_B^2} \right)^3 \zeta^2 [1 + \Delta R],$$

where ζ describes the flavor-SU(3) breaking between ρ/ω and K^* , and ΔR accounts for annihilation diagrams. Both ζ and ΔR must be taken from theory [3, 6, 23]. Following [3], we choose the values $\zeta = 0.85 \pm 0.10$, and $\Delta R = 0.10 \pm 0.10$, which is the average over the values given for the three modes. We find the limit $|V_{td}|/|V_{ts}| < 0.19$ at 90% C.L, ignoring the theoretical uncertainties. Our upper limit on $|V_{td}|/|V_{ts}|$ constrains $|V_{td}| < 0.008$ at 90% C.L. assuming $|V_{ts}| = |V_{cb}|$

[18]; this lies within the current 90% confidence interval $0.005 < |V_{td}| < 0.014$, which is obtained from a fit to experimental results on the CKM matrix elements [18]. Varying the values of ζ and ΔR within their uncertainties leads to changes in the limits by ± 0.03 and ± 0.001 for $|V_{td}|/|V_{ts}|$ and $|V_{td}|$, respectively.

In conclusion, we have found no evidence for the exclusive $b \rightarrow d\gamma$ transitions $B \rightarrow \rho\gamma$ and $B^0 \rightarrow \omega\gamma$ in 211 million $\Upsilon(4S) \rightarrow B\bar{B}$ decays studied with the BABAR detector. The 90% C.L. upper limits on the branching fractions and $|V_{td}|/|V_{ts}|$ are significantly lower than our previous values [7] and restrict the range indicated by SM predictions [3, 4].

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support BABAR. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONACyT (Mexico), A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

* Now at Department of Physics, University of Warwick, Coventry, United Kingdom

† Also with Università della Basilicata, Potenza, Italy

‡ Also with IFIC, Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Valencia, Spain

§ Deceased

[1] For a review, see K. Lingel, T. Skwarnicki, and J.G. Smith, *Ann. Rev. Nucl. Part. Sci.* **48**, 253 (1998).

[2] The named member of a charge-conjugate pair of parti-

cles stands for either.

- [3] A. Ali, E. Lunghi, and A. Parkhomenko, *Phys. Lett. B* **595**, 323 (2004).
- [4] S. W. Bosch and G. Buchalla, *Nucl. Phys. B* **621**, 459 (2002).
- [5] See, for example, S. Bertolini, F. Borzumati, and A. Masiero, *Nucl. Phys. B* **294**, 321 (1987); H. Baer and M. Brhlik, *Phys. Rev. D* **55**, 3201 (1997); J. Hewett and J. Wells, *Phys. Rev. D* **55**, 5549 (1997); M. Carena *et al.*, *Phys. Lett. B* **499**, 141 (2001).
- [6] A. Ali and A. Y. Parkhomenko, *Eur. Phys. J. C* **23**, 89 (2002).
- [7] B. Aubert *et al.* [BABAR Collaboration], *Phys. Rev. Lett.* **92**, 111801 (2004).
- [8] T.E. Coan *et al.* [CLEO Collaboration], *Phys. Rev. Lett.* **84**, 5283 (2000).
- [9] B. Aubert *et al.* [BABAR Collaboration], *Nucl. Instrum. Meth. A* **479**, 1 (2002).
- [10] B. Aubert *et al.* [BABAR Collaboration], *Phys. Rev. Lett.* **88**, 101805 (2002).
- [11] We use the Stuttgart Neural Network Simulator (<http://www-ra.informatik.uni-tuebingen.de/SNNS>) to train a neural net with one hidden layer of 5 or 6 nodes.
- [12] G.C. Fox and S. Wolfram, *Nucl. Phys. B* **149**, 413 (1979).
- [13] B. Aubert *et al.* [BABAR Collaboration], *Phys. Rev. Lett.* **89**, 201802 (2002).
- [14] R.A. Fisher, *Annals Eugen.* **7**, 179 (1936).
- [15] E.D. Bloom and C.W. Peck, *Ann. Rev. Nucl. Part. Sci.*, **33** 143 (1983).
- [16] H. Albrecht *et al.* [ARGUS Collaboration], *Z. Phys.* **C48**, 543 (1990).
- [17] K. Cranmer, *Comput. Phys. Commun.* **136**, 198 (2001).
- [18] S. Eidelman *et al.* [Particle Data Group Collaboration], *Phys. Lett. B* **592**, 1 (2004).
- [19] B. Aubert *et al.* [BABAR Collaboration], hep-ex/0403046; B. Aubert *et al.* [BABAR Collaboration], hep-ex/0403025.
- [20] B. Aubert *et al.* [BABAR Collaboration], *Phys. Rev. Lett.* **93**, 051802 (2004).
- [21] J. Dragic *et al.* [Belle Collaboration], hep-ex/0405068.
- [22] B. Aubert *et al.* [BABAR Collaboration], hep-ex/0407003.
- [23] B. Grinstein and D. Pirjol, *Phys. Rev. D* **62**, 093002 (2000).